



KENYATTA UNIVERSITY

UNIVERSITY EXAMINATIONS 2011/2012

**SECOND SEMESTER EXAMINATION FOR THE DEGREE OF BACHELOR
OF SCIENCE**

SMA 363: TESTS OF HYPOTHESES 1

DATE: WEDNESDAY 28TH MARCH 2012

TIME: 2.00 P.M. – 4.00 P.M.

INSTRUCTIONS

Answer question ONE and any other TWO questions

Question One (30 marks)

a) Distinguish the following terms as used in hypothesis testing

i) Type I and type II errors (2 marks)

ii) Simple and composite hypotheses (2 marks)

b) A random variable X is believed to follow an exponential distribution i.e.

$$f(x) = \begin{cases} \lambda e^{-\lambda x}, & x > 0, \lambda > 0 \\ 0 & \text{otherwise} \end{cases}$$

In order to test the null hypothesis $\mu = 20$ against the alternative hypothesis

$\mu = 30$, where $\mu = \frac{1}{\lambda}$, a single value is observed from this distribution. If this

value is less than 40, H_0 is accepted, otherwise H_0 is rejected. Find the probabilities of type I error and type II error, hence obtain the power of the test.

(6 marks)

- c) A random sample of 31 observations from a normal population possessed a mean $\bar{X} = 90$ and a standard deviation $S = 7.1$. Test $H_0: \sigma^2 = 40$ against $H_1: \sigma^2 > 40$. Use $\alpha = 5\%$ (5 marks)
- d) Suppose X is a uniform random variable over the interval $(-a, a)$, we wish to test $H_0: a = 1$ against $H_1: a > 1$. Find the size of the test if we take one observation of X and reject H_0 if $|x| > 0.99$ (4 marks)
- e) Suppose you were testing $H_0: \lambda = 2$ against $\lambda_1 = 1$ where λ is the parameter of the poisson distribution i.e.
- $$f(x) = \begin{cases} e^{-\lambda} \frac{\lambda^x}{x!}, & 0, 1, 2, \dots \\ 0 & \text{otherwise} \end{cases}$$
- Obtain the best critical region of the test. (8 marks)
- f) State the Neyman - Pearson Lemma for testing a simple hypothesis against a simple hypothesis. (3 marks)

Question Two (20 marks)

- a) Let $x_1, x_2, \dots, x_n$ be a random sample of size n from a normal population with unknown mean, μ , and unknown variance, σ^2 . Obtain the likelihood ratio critical region for testing the hypothesis $H_0: \sigma^2 = \sigma_0^2$ against $H_1: \sigma^2 \neq \sigma_0^2$, where σ_0^2 is specified. (12 marks)
- b) A random sample of $n = 7$ observations from a normal population produced the following measurements: 4, 0, 6, 3, 3, 5, 7. Do the data provide sufficient evidence to indicate that $\sigma^2 < 1$? Test using $\alpha = 5\%$ (8 marks)

Question Three (20 marks)

- a) Let $x_1, x_2, \dots, x_m$ be a random sample from a normal population with mean μ_1 and variance σ_1^2 , and $y_1, y_2, \dots, y_n$ be another independent random sample from a normal population with mean μ_2 and variance σ_2^2 .

Derive the test statistic for testing the hypothesis $H_0: \sigma_1^2 = \sigma_2^2 = \sigma^2$ against

$H_1: \sigma_1^2 \neq \sigma_2^2$. Take α to be the level of significance. (1 mark)

- b) Independent random samples were selected from each of two normally distributed populations $n_1 = 6$ from population 1 and $n_2 = 4$ from population 2. The data are shown in the table below

Sample 1	Sample 2
3.1	2.3
4.3	1.4
1.2	3.7
1.7	8.9
0.7	
3.4	

- i) Test $H_0: \sigma_1^2 = \sigma_2^2$ against $H_1: \sigma_1^2 < \sigma_2^2$. Use $\alpha = 0.05$
- ii) Test $H_0: \sigma_1^2 = \sigma_2^2$ against $H_1: \sigma_1^2 \neq \sigma_2^2$. Use $\alpha = 0.05$ (9 marks)

Question Four (20 marks)

- a) Given that X is a random variable which is normally distributed with unknown mean μ and random variance $\sigma^2 = 3$ and $x_1, x_2, \dots, x_n$ is a random sample of size n drawn from X . Obtain the critical region for testing the hypothesis $H_0: \mu = \mu_0$ against $H_1: \mu > \mu_0$. Take the level of significance to be α . (12 marks)

- b) Suppose observations from a normal population with variance $\sigma^2 = 4$ are
 $6, 3, 1, 2, 4, 2, 7, 8, 3, 5, 4$ and 9 . Test the hypothesis
 $H_0 : \mu = 2.4$ against $H_1 : \mu = 2.3$ at $\alpha = 0.05$ level of significance.
 (8 marks)

Question Five (20 marks)

- a) Consider a simple regression model

$$y_i = \alpha + \beta x_i + e_i \text{ where } \alpha, \beta \text{ are constant and } e_i \sim N(0, \sigma^2)$$

Derive a test statistic for testing the hypothesis $H_0 : \beta = 0$ against $H_1 : \beta \neq 0$. Take the level of significance to be $\alpha\%$
 (12 marks)

- b) Test the hypothesis $H_0 : \beta = 0$ against $H_1 : \beta \neq 0$ at $\alpha = 5\%$ if the observations are $(1, 2), (3, 2), (2, 10), (8, 6)$
 (8 marks)


700 copies

Table 2 CUMULATIVE NORMAL DISTRIBUTION

$$\Phi(x) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt$$


x	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

x	1.282	1.645	1.960	2.326	2.576	3.090	3.291	3.891	4.417
$\Phi(x)$	.90	.95	.975	.99	.995	.999	.9995	.99995	.999995
$2[1 - \Phi(x)]$	.20	.10	.05	.02	.01	.002	.001	.0001	.00001

TABLE VI Critical Values of t 

ν	$t_{.100}$	$t_{.050}$	$t_{.025}$	$t_{.010}$	$t_{.005}$	$t_{.001}$	$t_{.0005}$
1	3.078	6.314	12.706	31.821	63.657	318.31	636.62
2	1.886	2.920	4.303	6.965	9.925	22.326	31.598
3	1.638	2.353	3.182	4.541	5.841	10.213	12.924
4	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	1.330	1.734	2.101	2.552	2.878	3.610	3.922
19	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	1.326	1.725	2.086	2.528	2.845	3.552	3.850
21	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	1.319	1.714	2.069	2.500	2.807	3.485	3.767
24	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	1.310	1.697	2.042	2.457	2.750	3.385	3.646
40	1.303	1.684	2.021	2.423	2.704	3.307	3.551
60	1.296	1.671	2.000	2.390	2.660	3.232	3.460
120	1.289	1.658	1.980	2.358	2.617	3.160	3.373
∞	1.282	1.645	1.960	2.326	2.576	3.090	3.291

Source: From E. S. Pearson and H. O. Hartley (eds.), *The Biometrika Tables for Statisticians*, Vol. 1, 3d ed., Biometrika, 1966.

TABLE VII Critical Values of χ^2 


Degrees of Freedom	$\chi^2_{.995}$	$\chi^2_{.990}$	$\chi^2_{.975}$	$\chi^2_{.950}$	$\chi^2_{.900}$
1	.0000393	.0001571	.0009821	.0039321	.0157908
2	.0100251	.0201007	.0506356	.102587	.210720
3	.0717212	.114832	.215795	.351846	.584375
4	.206990	.297110	.484419	.710721	1.063623
5	.411740	.554300	.831211	1.145476	1.61031
6	.675727	.872085	1.237347	1.63539	2.20413
7	.989265	1.239043	1.68987	2.16735	2.83311
8	1.344419	1.646482	2.17973	2.73264	3.48954
9	1.734926	2.087912	2.70039	3.32511	4.16816
10	2.15585	2.55821	3.24697	3.94030	4.86518
11	2.60321	3.05347	3.81575	4.57481	5.57779
12	3.07382	3.57056	4.40379	5.22603	6.30380
13	3.56503	4.10691	5.00874	5.89186	7.04150
14	4.07468	4.66043	5.62872	6.57063	7.78953
15	4.60094	5.22935	6.26214	7.26094	8.54675
16	5.14224	5.81221	6.90766	7.96164	9.31223
17	5.69724	6.40776	7.56418	8.67176	10.0852
18	6.26481	7.01491	8.23075	9.39046	10.8649
19	6.84398	7.63273	8.90655	10.1170	11.6515
20	7.43386	8.26040	9.59083	10.8508	12.4426
21	8.03366	8.89720	10.28293	11.5913	13.2396
22	8.64272	9.54249	10.9823	12.3380	14.0415
23	9.26042	10.19567	11.6885	13.0905	14.8479
24	9.88623	10.8564	12.4011	13.8484	15.6587
25	10.5197	11.5240	13.1197	14.6114	16.4734
26	11.1603	12.1981	13.8439	15.3791	17.2919
27	11.8076	12.8786	14.5733	16.1513	18.1138
28	12.4613	13.5648	15.3079	16.9279	18.9392
29	13.1211	14.2565	16.0471	17.7083	19.7677
30	13.7867	14.9535	16.7908	18.4926	20.5992
40	20.7065	22.1643	24.4331	26.5093	29.0505
50	27.9907	29.7067	32.3574	34.7642	37.6886
60	35.5346	37.4848	40.4817	43.1879	46.4589
70	43.2752	45.4418	48.7576	51.7393	55.3290
80	51.1720	53.5400	57.1532	60.3915	64.2778
90	59.1963	61.7541	65.6466	69.1260	73.2912
100	67.3276	70.0648	74.2219	77.9295	82.3581

Source: From C. M. Thompson, "Tables of the Percentage Points of the χ^2 -Distribution," *Biometrika*, 1941, 32, 188-189.

continued

TABLE VII Continued

Degrees of Freedom	$\chi^2_{.100}$	$\chi^2_{.050}$	$\chi^2_{.025}$	$\chi^2_{.010}$	$\chi^2_{.005}$
1	2.70554	3.84146	5.02389	6.63490	7.87944
2	4.60517	5.99147	7.37776	9.21034	10.5966
3	6.25139	7.81473	9.34840	11.3449	12.8381
4	7.77944	9.48773	11.1433	13.2767	14.8602
5	9.23635	11.0705	12.8325	15.0863	16.7496
6	10.6446	12.5916	14.4494	16.8119	18.5476
7	12.0170	14.0671	16.0128	18.4753	20.2777
8	13.3616	15.5073	17.5346	20.0902	21.9550
9	14.6837	16.9190	19.0228	21.6660	23.5893
10	15.9871	18.3070	20.4831	23.2093	25.1882
11	17.2750	19.6751	21.9200	24.7250	26.7569
12	18.5494	21.0261	23.3367	26.2170	28.2995
13	19.8119	22.3621	24.7356	27.6883	29.8194
14	21.0642	23.6848	26.1190	29.1413	31.3193
15	22.3072	24.9958	27.4884	30.5779	32.8013
16	23.5418	26.2962	28.8454	31.9999	34.2672
17	24.7690	27.5871	30.1910	33.4087	35.7185
18	25.9894	28.8693	31.5264	34.8053	37.1564
19	27.2036	30.1435	32.8523	36.1908	38.5822
20	28.4120	31.4104	34.1696	37.5662	39.9968
21	29.6151	32.6705	35.4789	38.9321	41.4010
22	30.8133	33.9244	36.7807	40.2894	42.7956
23	32.0069	35.1725	38.0757	41.6384	44.1813
24	33.1963	36.4151	39.3641	42.9798	45.5585
25	34.3816	37.6525	40.6465	44.3141	46.9278
26	35.5631	38.8852	41.9232	45.6417	48.2899
27	36.7412	40.1133	43.1944	46.9630	49.6449
28	37.9159	41.3372	44.4607	48.2782	50.9933
29	39.0875	42.5569	45.7222	49.5879	52.3356
30	40.2560	43.7729	46.9792	50.8922	53.6720
40	51.8050	55.7585	59.3417	63.6907	66.7659
50	63.1671	67.5048	71.4202	76.1539	79.4900
60	74.3970	79.0819	83.2976	88.3794	91.9517
70	85.5271	90.5312	95.0231	100.425	104.215
80	96.5782	101.879	106.629	112.329	116.321
90	107.565	113.145	118.136	124.116	128.299
100	118.498	124.342	129.561	135.807	140.169

TABLE IX Percentage Points of the F -distribution, $\alpha = .05$ 


		NUMERATOR DEGREES OF FREEDOM								
ν_2	ν_1	1	2	3	4	5	6	7	8	9
1	1	161.4	199.5	215.7	224.6	230.2	234.0	236.8	238.9	240.5
2	1	18.51	19.00	19.16	19.25	19.30	19.33	19.35	19.37	19.38
3	1	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81
4	1	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00
5	1	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77
6	1	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10
7	1	5.59	4.71	4.33	4.10	3.97	3.87	3.79	3.73	3.68
8	1	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39
9	1	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18
10	1	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02
11	1	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90
12	1	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80
13	1	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71
14	1	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65
15	1	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59
16	1	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54
17	1	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49
18	1	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46
19	1	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42
20	1	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39
21	1	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37
22	1	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34
23	1	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32
24	1	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30
25	1	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28
26	1	4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27
27	1	4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25
28	1	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24
29	1	4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22
30	1	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21
40	1	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12
60	1	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04
120	1	3.92	3.07	2.68	2.45	2.29	2.17	2.09	2.02	1.96
∞	1	3.84	3.00	2.60	2.37	2.21	2.10	2.01	1.94	1.88

continued

Source: From M. Merrington and C. M. Thompson, "Tables of Percentage Points of the Inverted Beta (F)-Distribution." *Biometrika*, 1943, 33, 73-88.

TABLE IX Continued

ν_1	NUMERATOR DEGREES OF FREEDOM									
	10	12	15	20	24	30	40	60	120	∞
ν_2										
1	241.9	243.9	245.9	248.0	249.1	250.1	251.1	252.2	253.3	254.3
2	19.40	19.41	19.43	19.45	19.45	19.46	19.47	19.48	19.49	19.50
3	8.79	8.74	8.70	8.66	8.64	8.62	8.59	8.57	8.55	8.53
4	5.96	5.91	5.86	5.80	5.77	5.75	5.72	5.69	5.66	5.63
5	4.74	4.68	4.62	4.56	4.53	4.50	4.46	4.43	4.40	4.36
6	4.06	4.00	3.94	3.87	3.84	3.81	3.77	3.74	3.70	3.67
7	3.64	3.57	3.51	3.44	3.41	3.38	3.34	3.30	3.27	3.23
8	3.35	3.28	3.22	3.15	3.12	3.08	3.04	3.01	2.97	2.93
9	3.14	3.07	3.01	2.94	2.90	2.86	2.83	2.79	2.75	2.71
10	2.98	2.91	2.85	2.77	2.74	2.70	2.66	2.62	2.58	2.54
11	2.85	2.79	2.72	2.65	2.61	2.57	2.53	2.49	2.45	2.40
12	2.75	2.69	2.62	2.54	2.51	2.47	2.43	2.38	2.34	2.30
13	2.67	2.60	2.53	2.46	2.42	2.38	2.34	2.30	2.25	2.21
14	2.60	2.53	2.46	2.39	2.35	2.31	2.27	2.22	2.18	2.13
15	2.54	2.48	2.40	2.33	2.29	2.25	2.20	2.16	2.11	2.07
16	2.49	2.42	2.35	2.28	2.24	2.19	2.15	2.11	2.06	2.01
17	2.45	2.38	2.31	2.23	2.19	2.15	2.10	2.06	2.01	1.96
18	2.41	2.34	2.27	2.19	2.15	2.11	2.06	2.02	1.97	1.92
19	2.38	2.31	2.23	2.16	2.11	2.07	2.03	1.98	1.93	1.88
20	2.35	2.28	2.20	2.12	2.08	2.04	1.99	1.95	1.90	1.84
21	2.32	2.25	2.18	2.10	2.05	2.01	1.96	1.92	1.87	1.81
22	2.30	2.23	2.15	2.07	2.03	1.98	1.94	1.89	1.84	1.78
23	2.27	2.20	2.13	2.05	2.01	1.96	1.91	1.86	1.81	1.76
24	2.25	2.18	2.11	2.03	1.98	1.94	1.89	1.84	1.79	1.73
25	2.24	2.16	2.09	2.01	1.96	1.92	1.87	1.82	1.77	1.71
26	2.22	2.15	2.07	1.99	1.95	1.90	1.85	1.80	1.75	1.69
27	2.20	2.13	2.06	1.97	1.93	1.88	1.84	1.79	1.73	1.67
28	2.19	2.12	2.04	1.96	1.91	1.87	1.82	1.77	1.71	1.65
29	2.18	2.10	2.03	1.94	1.90	1.85	1.81	1.75	1.70	1.64
30	2.16	2.09	2.01	1.93	1.89	1.84	1.79	1.74	1.68	1.62
40	2.08	2.00	1.92	1.84	1.79	1.74	1.69	1.64	1.58	1.51
60	1.99	1.92	1.84	1.75	1.70	1.65	1.59	1.53	1.47	1.39
120	1.91	1.83	1.75	1.66	1.61	1.55	1.50	1.43	1.35	1.25
∞	1.83	1.75	1.67	1.57	1.52	1.46	1.39	1.32	1.22	1.00

TABLE X Percentage Points of the F -distribution, $\alpha = .025$ 

		NUMERATOR DEGREES OF FREEDOM								
ν_1	ν_2	1	2	3	4	5	6	7	8	9
	1	647.8	799.5	864.2	899.6	921.8	937.1	948.2	956.7	963.3
	2	38.51	39.00	39.17	39.25	39.30	39.33	39.36	39.37	39.39
	3	17.44	16.04	15.44	15.10	14.88	14.73	14.62	14.54	14.47
	4	12.22	10.65	9.98	9.60	9.36	9.20	9.07	8.98	8.90
	5	10.01	8.43	7.76	7.39	7.15	6.98	6.85	6.76	6.68
	6	8.81	7.26	6.60	6.23	5.99	5.82	5.70	5.60	5.52
	7	8.07	6.54	5.89	5.52	5.29	5.12	4.99	4.90	4.82
	8	7.57	6.06	5.42	5.05	4.82	4.65	4.53	4.43	4.36
	9	7.21	5.71	5.08	4.72	4.48	4.32	4.20	4.10	4.03
Denominator Degrees of Freedom	11	6.72	5.26	4.63	4.28	4.04	3.88	3.76	3.66	3.59
	12	6.55	5.10	4.47	4.12	3.89	3.73	3.61	3.51	3.44
	13	6.41	4.97	4.35	4.00	3.77	3.60	3.48	3.39	3.31
	14	6.30	4.86	4.24	3.89	3.66	3.50	3.38	3.29	3.21
	15	6.20	4.77	4.15	3.80	3.58	3.41	3.29	3.20	3.12
	16	6.12	4.69	4.08	3.73	3.50	3.34	3.22	3.12	3.05
	17	6.04	4.62	4.01	3.66	3.44	3.28	3.16	3.06	2.98
	18	5.98	4.56	3.95	3.61	3.38	3.22	3.10	3.01	2.93
	19	5.92	4.51	3.90	3.56	3.33	3.17	3.05	2.96	2.88
	20	5.87	4.46	3.86	3.51	3.29	3.13	3.01	2.91	2.84
	21	5.83	4.42	3.82	3.48	3.25	3.09	2.97	2.87	2.80
	22	5.79	4.38	3.78	3.44	3.22	3.05	2.93	2.84	2.76
	23	5.75	4.35	3.75	3.41	3.18	3.02	2.90	2.81	2.73
	24	5.72	4.32	3.72	3.38	3.15	2.99	2.87	2.78	2.70
	25	5.69	4.29	3.69	3.35	3.13	2.97	2.85	2.75	2.68
	26	5.66	4.27	3.67	3.33	3.10	2.94	2.82	2.73	2.65
	27	5.63	4.24	3.65	3.31	3.08	2.92	2.80	2.71	2.63
	28	5.61	4.22	3.63	3.29	3.06	2.90	2.78	2.69	2.61
	29	5.59	4.20	3.61	3.27	3.04	2.88	2.76	2.67	2.59
	30	5.57	4.18	3.59	3.25	3.03	2.87	2.75	2.65	2.57
	40	5.42	4.05	3.46	3.13	2.90	2.74	2.62	2.53	2.45
	60	5.29	3.93	3.34	3.01	2.79	2.63	2.51	2.41	2.33
	120	5.15	3.80	3.23	2.89	2.67	2.52	2.39	2.30	2.22
	∞	5.02	3.69	3.12	2.79	2.57	2.41	2.29	2.19	2.11

continued

Source: From M. Merrington and C. M. Thompson, "Tables of Percentage Points of the Inverted Beta (F)-Distribution," *Biometrika*, 1943, 33, 73-88.